

Critical numbers of attractive Bose-condensed atoms in asymmetric traps

A. Gammal^(a), Lauro Tomio^(b) and T.Frederico^(c)

^(a) *Instituto de Física, Universidade de São Paulo, 05315-970 São Paulo, Brazil*

^(b) *Instituto de Física Teórica, Universidade Estadual Paulista, 01405-900 São Paulo, Brazil*

^(c) *Departamento de Física, Instituto Tecnológico da Aeronáutica, CTA, 12228-900 São José dos Campos, SP, Brazil*

(February 1, 2008)

The recent Bose-Einstein condensation of ultracold atoms with attractive interactions led us to consider the novel possibility to probe the stability of its ground state in arbitrary three-dimensional harmonic traps. We performed a quantitative analysis of the critical number of atoms through a full numerical solution of the mean field Gross-Pitaevskii equation. Characteristic limits are obtained for reductions from three to two and one dimensions, in perfect cylindrical symmetries as well as in deformed ones.

PACS: 03.75.Fi, 32.80.Pj, 42.65.Tg, 02.60.Jh

The predicted collapsing behavior of the condensed systems with attractive two-body atomic interactions [1], first observed in experiments with ^7Li [2], was recently tested in experiments with ^{85}Rb [3]. In the experiments described in Ref. [3], and more recently in [4], by means of Feshbach resonance techniques [5], the two-body interaction was tuned from positive to negative values. Besides the fact that the experimental results qualitatively agree with the theory, and confirm results of previous variational treatments [6,7,8,9,10,11], they also show a consistent quantitative deviation of about 20% from the mean-field predicted critical number of atoms, N_c [1]. The asymmetry of the trap was shown in Ref. [12] to be responsible for about 4% of the observed deviation.

In this respect, it is relevant to obtain precise and reliable numerical results for the mean-field calculations, in order to probe their consistency and possible limitations. The actual experimental atomic traps are in general harmonic and non-symmetric. Extreme asymmetric traps have been recently employed in experimental investigations with condensates constrained to quasi-one (1D) [4,13] or quasi-two dimensions (2D) [14], exploring theoretical analysis considered by several authors [15,16,17,7,18,19]. A non symmetric three-dimensional (3D) trap is reported in Ref. [20], with the frequencies given by $2\omega_1 = \sqrt{2}\omega_2 = \omega_3 = 2\pi \times 33\text{Hz}$.

Considering the general non-symmetric traps that have been employed, the accuracy of the comparison between experiments and the results of mean-field approximation relies in precise calculations using arbitrary three-dimensional traps. In case of attractive two-body interaction, the maximum critical number of atoms for a stable system is one of the interesting observables to study, which is also related to the collapse of the wave-function of the system. In these cases, where the two-body scattering length is negative and the kinetic energy cannot be considered to be a small perturbation, the Gross-Pitaevskii mean field approximation has been applied, given reliable results in explaining the observations in the stable (non-collapsing) conditions [21].

Before presenting the mean-field equation for an arbitrary

3D case, let us analyze qualitatively the collapse phenomenon for asymmetric traps. The interaction energy is proportional to the square of the density, varying with the negative two-body scattering length. For traps with cylindrical (or almost cylindrical) shapes, there are two quite different situations: one, pancake-like, with the frequencies in the transverse directions being smaller than the frequency in the longitudinal direction; the other, cigar-like (quasi-1D), with the frequency in the longitudinal direction smaller than the frequencies in the perpendicular directions. For a true 1D system, one does not expect the collapse of the system with increasing number of atoms [1,22]. However, it happens that a realistic 1D limit is not a true 1D system, with the density of particles still increasing due to the strong restoring forces in the perpendicular directions [16]. The relevance of the quasi-1D trap have been pointed out in Ref. [16], to control the condensate motion. But, as we are going to see, the critical number of particles in the quasi-1D limit is smaller than the critical number of particles in the 2D limit, if we just exchange the longitudinal and perpendicular frequencies. The physical reason for that behavior is the increase of the average density in the cigar-like configuration relative to the pancake like one for the same number of atoms, implying in a strong collapsing force in the first case and consequently the cigar-like geometry is a more unstable configuration compared to the pancake like one. This conclusion is in apparent contradiction with the remark made in section IV of Ref. [16], saying that, considering the better collapse-avoiding properties, “the cigar-shaped trap is the optimal one”. We are going to discuss this problem in detail and clarify this issue.

In the following, we revise the Gross-Pitaevskii (GP) formalism, for an atomic system with arbitrary non-spherically symmetric harmonic trap. The Bose-Einstein condensate, at zero temperature, in the GP mean-field approximation is given by

$$i\hbar \frac{\partial}{\partial t} \Psi(\vec{r}, t) = \left[-\frac{\hbar^2}{2m} \nabla^2 + \frac{m}{2} (\omega_1^2 r_1^2 + \omega_2^2 r_2^2 + \omega_3^2 r_3^2) \right] \Psi(\vec{r}, t)$$

$$+ \frac{4\pi\hbar^2 a}{m} |\Psi(\vec{r}, t)|^2 \Big] \Psi(\vec{r}, t) = \mu \Psi(\vec{r}, t), \quad (1)$$

where $r = \sqrt{r_1^2 + r_2^2 + r_3^2}$, m is the mass of the atom, μ is the chemical potential, and the wave-function $\Psi \equiv \Psi(\vec{r}, t) = \Psi(\vec{r}, 0) \exp(-i\mu t/\hbar)$ is normalized to the number of particles N . The arbitrary geometry of the trap is parameterized by three different frequencies ω_1 , ω_2 and ω_3 . For convenience, it is appropriate to define the frequencies according to their magnitude, such that, in the present work we assume $\omega_1 \leq \omega_2 \leq \omega_3$.

Here we will be concerned only with systems that have attractive two-body interactions [$a = -|a|$, in Eq. (1)]. In this case, it was first shown numerically, in Ref. [1], that the system becomes unstable if a maximum critical number of atoms, N_c , is achieved. We present precise results for a critical parameter k , directly related to the maximum number of atoms, in a general non-symmetric configuration of the trap.

By rewriting Eq.(1) in dimensionless units:

$$i \frac{\partial \phi}{\partial \tau} = \left[\frac{1}{2} \sum_i \left(-\frac{\partial^2}{\partial x_i^2} + \frac{\omega_i^2 x_i^2}{\omega^2} \right) - |\phi|^2 \right] \phi, \quad (2)$$

where $\tau \equiv \omega t$, $r_i \equiv l_0 x_i$ and $\phi \equiv l_0 \sqrt{4\pi|a|} \Psi$, with

$$\int d^3x |\phi|^2 = 4\pi \frac{N|a|}{l_0} \quad (3)$$

The oscillator length l_0 is defined in terms of ω , which is taken as the geometrical mean value of the frequencies:

$$l_0 \equiv \sqrt{\frac{\hbar}{m\omega}}, \quad \text{with } \omega \equiv (\omega_1 \omega_2 \omega_3)^{1/3}. \quad (4)$$

For strong non-symmetric cases, particularly when comparing the two extreme cylindrical-shape geometries, $\omega_1 \sim \omega_2 \ll \omega_3$ (pancake-shape) and $\omega_1 \ll \omega_2 \sim \omega_3$ (cigar-shape), it is expected a noticeable difference between the corresponding critical number of particles.

We define a parameter k , related to the critical number of trapped atoms N_c as in Ref. [3]:

$$k = \frac{N_c |a|}{l_0}. \quad (5)$$

This parameter is a maximum critical limit for stable solutions of the dimensionless Eq. (2). It will depend only on the ratio of the frequencies of the trap. Within the precision given in Ref. [23], $k_s = 0.5746$, where k_s is k for spherically symmetric traps. In Ref. [24], the critical number was calculated for a nonsymmetrical geometry, but in a case that the frequency ratio is not too far from the unity ($\omega_1/\omega_\perp = 0.72$), giving a result for the number of atoms almost equal to the spherical one.

In the experiments with ^{85}Rb [3], they have considered an almost cylindrical “cigar-type” symmetry, with the three frequencies given by 17.47 Hz, 17.24 Hz and 6.80

Hz. With this symmetry, they have obtained $k = 0.459 \pm 0.012$ (statistical) ± 0.054 (systematic). In Ref. [12], assuming the cylindrical symmetry $\omega_1 = 2\pi \times 6.80\text{Hz}$, $\omega_2 = \omega_3 = 2\pi \times 17.35\text{ Hz}$, it was obtained $k = 0.55$, a value about 4% lower than k_s .

In our numerical approach, the calculation is performed by evolving the nonlinear equation (2) through imaginary time [24]. The evolution is performed for an initial value of the normalization (3) till the wave function relaxes to the ground state. The wave function is renormalized after each time step. The process is repeated systematically for larger values of the normalization, until a critical limit is reached. At this critical limit the ground state becomes unstable. The time evolution is done with a semi-implicit second order finite difference algorithm. An alternating scheme is used in the x_1 and x_2 direction, with a split step in the x_3 direction. This procedure is done only for $x_i \geq 0$, taking advantage of the reflection symmetry of the ground state. We consider 100^3 grid points and time step equal to $\Delta\tau = 0.001$, verifying that the algorithm is stable for long time evolution. As we increment the normalization, approaching the critical limit, the wave function starts to shrink. So, in order to maintain the precision, we introduce an automatic reduction of the grid sizes, Δx_1 , Δx_2 , and Δx_3 , gauged by the respective root mean-square-radius in each direction.

In Fig. 1 we show our main results for the critical constant k , covering many different geometries. We plot k/k_s (and k in the rhs y -axis) as a function of λ , which is defined by

$$\lambda \equiv \frac{\omega_1 \omega_3}{\omega_2^2} = \left(\frac{\omega}{\omega_2} \right)^3, \quad \text{with } \omega_3 \geq \omega_2 \geq \omega_1. \quad (6)$$




FIG. 1. Critical constant $k = N_c |a| (m\omega/\hbar)^{1/2}$, calculated for an arbitrary geometry with $\omega_3 \geq \omega_2 \geq \omega_1$. The ratio ω_3/ω_1 is shown below each corresponding curve. The dotted-lines correspond to cylindrical ($\eta = 1$) and deformed cylindrical ($\eta = 1.414$) geometries.

We show several curves in which we kept constant the ratio ω_3/ω_1 (solid-lines). The values of ω_3/ω_1 are indicated inside the figure, just below the corresponding plot. The dashed and dotted-lines correspond to cylindrical ($\eta = 1$) and deformed cylindrical ($\eta = 1.414$) symmetries for the trap, where η is the deformation parameter, as it will be explained. In the left-hand-side (lhs) of these plots we have $\omega_3 = \eta\omega_2 \gg \omega_1$ (cigar-shape); and, in the right-hand-side, $\omega_3 \gg \omega_2 = \eta\omega_1$ (pancake-shape). k can be determined for any symmetry that was not shown, by interpolating the already given results. The results for the complete non-symmetric case are consistent with the previously obtained values in cylindrical symmetry [12]. The maximum value for the critical number k is obtained for the spherically symmetric case ($\omega_1 = \omega_2 = \omega_3$).

As already observed, Fig. 1 includes previous calculations in the limit of the quasi 1D (cigar-shape) and quasi 2D (pancake-shape) symmetries (see dashed-line). However, for the sake of comparison with previous results obtained by several authors, we also present in Fig. 2 the cylindrical pancake-type ($\omega_\perp = \omega_1 = \omega_2 \ll \omega_3$) and cigar-type ($\omega_\perp = \omega_3 = \omega_2 \gg \omega_1$) results. In Fig. 2, for cylindrical geometries, we compare our exact results, for $k_3 = N_c|a|(m\omega_3/\hbar)^{1/2}$ as a function of λ , with the corresponding variational ones of Refs. [7,15]. The variational results (dashed-lines) are consistently a bit higher than the exact ones. For $\lambda \rightarrow 0$ ($\omega_1 \ll \omega_2 = \omega_3 = \omega_\perp$), the exact and variational results for the critical constant are $N_c|a|(m\omega_\perp/\hbar)^{1/2} = 0.676$ and 0.776 , respectively. They are consistent with the quasi-1D limits given in Refs. [15,16]. When $\omega_3 \gg \omega_2 = \omega_1$, the variational 2D limit $\sqrt{\pi/2}$ of Ref. [7] is comparable with our exact result ($k_3 = 0.931\sqrt{\pi/2}$). In this case, the quasi-2D limit coincide with the true 2D limit [25,19,22,26].




FIG. 2. Exact results of $k_3 \equiv N_c|a|(m\omega_3/\hbar)^{1/2}$, for cylindrical traps (solid-lines) compared with variational approaches (dashed-lines). In the 2D pancake limit, $\omega_\perp = \omega_1 = \omega_2 = \omega_3$ and $k_3 = k\lambda^{-1/6}$. Note that $\lambda = \omega_3/\omega_1$, when $\omega_2 = \omega_3$.

Considering the present analysis, we observe that N_c in a cigar-like (quasi-1D) trap is smaller than N_c in a pancake-like (quasi-2D) trap. And, as we deform a cigar-

like trap, N_c also increases. We need to clarify this matter, which is the subject of the next four paragraphs, because such result is apparently contradicting a remark made in Ref. [16], saying that “the cigar-shaped trap is the optimal one”.

As shown in Ref. [12] and also in the present calculation, in a deformed cylindrical symmetry, the cigar shape (with one of the frequencies smaller than the other two) is more favorable to obtain a larger value of k than the pancake-shape symmetry (with one of the frequencies larger than the other two). See, for example, in Fig. 1, the two extreme points of the curve with $\omega_3/\omega_1 = 100$. This results from the definition of k , Eq. (5), in terms of the averaged oscillator length l_0 . However, the maximum value of k can only be directly related with the maximum value of N_c in case that ω is kept fixed. And, with l_0 fixed, N_c is maximized for $\lambda = 1$, corresponding to the spherically symmetric case ($k = k_s$). If we fix l_0 and ω_3/ω_1 , N_c is maximized for a deformed cylindrical symmetry with $\lambda < 1$, as one can see from Fig. 1.

Considering exact cylindrical traps, by exchanging the frequencies (which, obviously, does not keep constant the averaged frequency ω), it is valid the following ratio that was obtained in Ref. [12]:

$$R(\lambda) \equiv \frac{N_c(\lambda)}{N_c(1/\lambda)} = \lambda^{1/6} \frac{k(\lambda)}{k(1/\lambda)}. \quad (7)$$

This result favors the pancake-like symmetry ($\lambda = \omega_3/\omega_1 = \omega_3/\omega_\perp > 1$), to obtain a larger value for N_c . Consider, for example, a cylindrical pancake-type trap with $\lambda = 100$, in comparison with a cigar-type trap with $\lambda = 1/100$. We notice that, in this case, $R(\lambda = 100) \approx 1.6$, implying that with such pancake-like trap ($\omega_3 = 100\omega_\perp$) one can obtain about 60% more particles than with the corresponding cigar-like trap ($\omega_3 = \omega_\perp = 100\omega_1$). Let us consider the recent experiment with quasi-1D (cigar-like) trap used in the formation and propagation of matter wave solitons, with ^7Li [4]. In this case, it was used axial and radial frequencies, respectively, equal to 3.2 Hz and ~ 625 Hz [29]; or, $\omega_\perp = \omega_3 = 2\pi \times 625$ Hz, $\omega_1 = 2\pi \times 3.2$ Hz and $\lambda = \omega_1/\omega_3 = 0.00513$. So, as shown in Fig. 2, we are practically in the limit $\lambda \rightarrow 0$, which gives $N_c^{1D}|a|/\sqrt{\hbar/(m\omega_\perp)} \approx 0.675$. Considering that the scattering length was tuned to $a = -3a_0$ (a_0 is the Bohr radius), the maximum number of atoms in this quasi-1D trap is $N_c^{1D} \approx 6400$. If we exchange the radial and axial frequencies in this experiment, going from a cigar-like to a pancake-like trap, $\omega_\perp = 2\pi \times 3.2$ Hz, $\omega_3 = 2\pi \times 625$ Hz and $\lambda = \omega_3/\omega_\perp = 195.3$. In this case, $N_c^{2D}|a|/\sqrt{\hbar/(m\omega_3)} \approx 1.12$. So, the critical number in this quasi-2D trap is $N_c^{2D} \approx 10600$, which is about 66% larger than the corresponding number of atoms in the quasi-1D case, $N_c^{1D} \approx 6400$.

The discussion about the best way to distribute the frequencies to obtain the maximum number of atoms was first considered in Ref. [6], arriving that the best is to do a spherical trap, for achieving maximum density. (In

their almost spherical trap they have obtained from ~ 600 to ~ 1300 atoms, in an overall agreement with theoretical predictions [2].) The point is that, to say what is the best configuration for the maximum number of atoms we must also say what is the *constraint* in the frequencies. If the constraint is to keep fixed the product $\omega_1\omega_2\omega_3$ than the maximum number will be given for the maximum k [as shown in Eq.(5)], that happens in a spherical symmetry, in agreement with Ref. [6]. But, if two frequencies are equal and given *a priori*, than the best configuration to increase N_c is to make the third frequency equal to zero, in agreement with Ref. [16]. However, if only one frequency is kept fixed, than the extreme pancake shape will contain more atoms, by making the remaining frequencies go to zero.

As a final remark about the maximization of N_c , we note that, for attractive two-body interactions, the frequencies in all three directions cannot be initially zero. In the experimental process of condensation, the atoms must be trapped while going from higher temperature T to the critical one, T_c . By making the frequencies too small, the T_c will be smaller and the condensation harder to achieve [as shown in Ref. [27], $T_c \sim \hbar\omega(N_{tot})^{1/3}$, where N_{tot} is the total number of particles in the trap]. Once the condensate is achieved, we can switch off some of the frequencies, and obtain a finite maximum value for N_c , as observed in Fig. 2 for $T = 0$. In the present work, we are concerned with the case of zero temperature. Effects of temperature on the collapse and on N_c , for spherically symmetric case, have been discussed, for example, in Ref. [28]. N_c is maximized by making ω as small as possible. However, a discontinuity exists when taking all the three frequencies exactly zero. Without trap the collapse will occur for any number of particles.

We must stress that Fig. 1 also includes novel non-symmetric cigar-type geometries $\omega_3 = \eta\omega_2 \gg \omega_1$ and elliptical pancake-type geometries $\omega_3 \gg \omega_2 = \eta\omega_1$, where η is the excentricity of the ellipsis. In particular, the dotted line in Fig.1 gives the critical values of k for $\eta = \sqrt{2}$. Thus, one could replot, as in Fig. 2, a series of curves whose limiting cases describe non-symmetric cigar- or pancake-like symmetries. In the 1D and 2D limits, one can obtain critical quasi-1D and quasi-2D numbers, for each η .

One can also obtain an interesting result, for asymmetric cigar-shape traps, where $\omega_3 = \eta\omega_2 \gg \omega_1$, following the approach given in section II of Ref. [16]. The assumption made in [16] to reduce the 3D equation to a 1D equation can be justified in the limit when the forces in the transverse directions are the main responsible for the trapping potential. So, in the limit $\omega_1 \rightarrow 0$, we are able to generalize the solitonic 1D equation for a cigar-shaped trap, deformed by a given ratio $\eta = \omega_3/\omega_2$ between the transverse direction frequencies. In this case, $\eta \geq 1$ cannot be arbitrarily larger. Also, by a scaling procedure applied to the equation for different values of η , using our previous exact result, we have

$$N_c|a| \left(\frac{m\sqrt{\omega_2\omega_3}}{\hbar} \right)^{1/2} = \frac{N_c|a|}{\eta^{1/4}} \sqrt{\frac{m\omega_3}{\hbar}} = 0.676. \quad (8)$$

This generalizes the cigar-shape quasi-1D results of [16] (with η such that we still have $\omega_2 \gg \omega_1$). The 1D solid line given in Fig.2 is also applied to deformed cigar-shaped symmetries if we replace k_3 by $k_3\eta^{-1/4}$ in the y -axis. This result may be relevant for asymmetric waveguide propagation as one can deform the cigar-type symmetry and control the collapsing condition. From Eq. (8) we observe that the maximum critical number N_c will increase when deforming the cigar-like symmetry by a factor proportional to $(\omega_3/\omega_2)^{1/4}$.

Now, let us see the effect of deformation in the same example of the cigar-shaped geometry used in Ref. [4], with ^7Li gas. We use the same value of $a = -3a_0$, with $\omega_3 = 2\pi \times 625 \text{ s}^{-1}$, and take $\omega_2 = \omega_3/\eta$. As given in Eq. (8), we go from $N_c = 6392$ ($\eta = 1$) to $N_c = 6970$ ($\eta = \sqrt{2}$), or $N_c = 7601$ ($\eta = 2$). It means, an increase of $\sim 9\%$ when $\eta = \sqrt{2}$; and $\sim 19\%$ when $\eta = 2$. In case, $\eta \gg 1$, the approximation considered in the wavefunction separation, as given in [16], is not valid. In such case we are reaching the other deformed pancake-like symmetry, where $\omega_3 \gg \omega_2 \sim \omega_1$. However, in the pancake geometry, the effect of deformation in N_c is negligible. By comparing the quasi pancake-like geometry with the quasi cigar-like geometry, the number N_c in the cigar-like geometry is much more sensitive to deformations.




FIG. 3. The three components, $i = 1, 2, 3$, of the mean-square-radius, $\langle x_i^2 \rangle$, $N|a|/l_0$, for the case that $\omega_3 = \sqrt{2}\omega$, $\omega_2 = \omega$, and $\omega_1 = \omega/\sqrt{2}$.

We have also studied the behavior of the root mean square radius for the case $\omega_3 = \sqrt{2}\omega_2 = 2\omega_1$. We verified that, as the system approaches the critical point (or collapse), the wave function tends to be more “spherical”,

confirming earlier conclusion made with a Gaussian variational approximation [30]. In Fig. 3, we show the corresponding results, for the three components of the mean-square-radius, $\langle r_i^2 \rangle = l_0^2 \langle x_i^2 \rangle$ ($i = 1, 2, 3$). As shown, when $N = 0$, we have $\langle r_1^2 \rangle / \langle r_2^2 \rangle = \langle r_2^2 \rangle / \langle r_3^2 \rangle = \sqrt{2}$; and, when $N \approx N_c$ such ratio is drastically reduced.

In summary, we have calculated systematically the critical number of particles, in systems that have negative two-body interactions, for traps with arbitrary geometries. The maximum critical number of particles N_c can be derived from the given value of the parameter k , given in Eq. (5), once one has the scattering length, the mass of the atomic system, and the frequencies of the trap. The results are shown in Figs. 1 and 2. The value of k , for any symmetry non explicitly given, can be easily derived from Fig. 1, by interpolation. It is also pointed out that the results shown in Fig. 2, in the 1D cigar-like case, can be extended for slightly deformed cylindrical symmetries, by replacing the y -axis label k_3 by $k_3 \eta^{-1/4}$.

Our main results in the present work are: (i) The maximum number of particles, N_c , for arbitrary 3D trap geometries is given through the results shown in Fig.1. (ii) The optimal trap configuration, to avoid the collapse with maximum N_c , is found to be strongly dependent on the constraints of the frequencies of the trap. If we initially fix one of the frequencies, the best configuration of the trap is pancake-like, with the other two frequencies going to zero. Analogously, if we initially fix two equal frequencies, the best configuration of the trap is cigar-like, with the third frequency close to zero. If we initially fix two different frequencies and try to vary the third frequency between the fixed ones, the best configuration is again pancake-like. We show that N_c is much more sensitive to deformations of the trap in a cigar-like geometry than in a pancake-like geometry. Finally, for small deformations η of the cigar-like traps, where $\eta \geq 1$, releasing the longitudinal direction, the solitonic solutions, obtained in Ref. [16] will be rescaled by the deformation. N_c will be rescaled by a factor $\eta^{1/4}$, generalizing the findings of Ref. [16].

ACKNOWLEDGEMENTS

We are grateful to Randy Hulet for informations provided. AG also thanks Emerson Passos and Marcelo Pires for useful discussions. For partial support, we thank Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

- [1] P.A. Ruprecht, M.J. Holland, K. Burnett, and M. Edwards, Phys. Rev. A **51**, 4704 (1995).
- [2] C.C. Bradley, C.A. Sackett, J.J. Tollett, and R.G. Hulet, Phys. Rev. Lett. **75**, 1687 (1995); C.C. Bradley, C.A. Sackett and R.G. Hulet, Phys. Rev. Lett. **78**, 985 (1997); J. M. Gerton, D. Strekalov, I. Prodan, and R. G. Hulet, Nature **408**, 692 (2000).
- [3] J.L. Roberts *et al.*, Phys. Rev. Lett. **86**, 4211 (2001); E. Donley *et al.*, Nature **412**, 295 (2001).
- [4] K.E. Strecker, G.B. Partridge, A.G. Truscott, and R.G. Hulet, Nature **417**, 150 (2002).
- [5] S. Inouye *et al.*, Nature **392**, 151 (1998); Ph. Courteille, R.S. Freeland, D. J. Heinzen, F.A. van Abeelen, and B.J. Verhaar, Phys. Rev. Lett **81**, 69 (1998).
- [6] C.A. Sackett, C.C. Bradley, M. Welling, and R.G. Hulet, Appl. Phys. B **65**, 433 (1997).
- [7] G. Baym and C.J. Pethick, Phys. Rev. Lett. **76**, 6 (1996).
- [8] E.V. Shuryak, Phys. Rev. A **54**, 3151 (1996).
- [9] H.T.C. Stoof, J. Stat. Phys. **87**, 1353 (1997).
- [10] M.Ueda and A.J.Leggett, Phys. Rev. Lett. **80**, 1576 (1998).
- [11] M. Wadati and T. Tsurumi, J. Phys. Soc. Japan **68**, 3840 (1999).
- [12] A. Gammal, T. Frederico, and L. Tomio, Phys. Rev. A **64**, 055602 (2001).
- [13] F. Schreck *et al.*, Phys. Rev. Lett. **87**, 080403 (2001); F. Schreck *et al.*, Phys. Rev. A **64**, 011402 (2001).
- [14] A. Görlitz *et al.*, Phys. Rev. Lett. **87**, 130402 (2001).
- [15] V.M. Pérez-Garcia, H. Michinel, J.I. Cirac, M. Lewenstein, and P. Zoller, Phys. Rev. A **56**, 1424 (1997).
- [16] V.M. Pérez-Garcia, H. Michinel, and H. Herrero, Phys. Rev. A **57**, 3837 (1998).
- [17] L.D. Carr, C.W. Clark, and W.P. Reinhardt, Phys. Rev. A **62**, 063610 (2000); Phys. Rev. A **62**, 063611 (2000); L.D. Carr, M.A. Leung, and W.P. Reinhardt, J. of Phys. B **33**, 3983 (2000).
- [18] R.K. Bhaduri, S.M. Reimann, S. Viefers, A. G. Choudhury, and M.K. Srivastava, J. of Phys. B **33**, 3895 (2000).
- [19] S.K. Adhikari, Phys. Rev. E **62**, 2937 (2000).
- [20] W.K. Hensinger *et al.*, Nature **412**, 52 (2001).
- [21] J.R. Anglin and W. Ketterle, Nature **416**, 211 (2002).
- [22] A. Gammal, T. Frederico, L. Tomio, F.K. Abdullaev, Phys. Lett. A **267**, 305 (2000).
- [23] A. Gammal, T. Frederico, L. Tomio, Phys. Rev. E **60**, 2421 (1999).
- [24] F. Dalfovo and S. Stringari, Phys. Rev. A **53**, 2477 (1996).
- [25] M.I. Weinstein, Commun. Math. Phys. **87**, 567 (1983).
- [26] L. Bergé, T.J. Alexander, and Y.S. Kivshar, Phys. Rev. A **62**, 023607 (2000).
- [27] F. Dalfovo, S. Giorgini, L.P. Pitaevskii, and S. Stringari, Rev. of Mod. Phys. **71**, 463 (1999).
- [28] M.J. Davis, D.A.W. Hutchinson, and E. Zaremba, J. Phys. B **32**, 3993 (1999).
- [29] R.G. Hulet, *private communication*.
- [30] M.O.D. Pires and E.J.V. de Passos, J. Phys. B.: At. Mol. Opt., **33**, 3929 (2000).